arXiv:1608.05925v1 [math.NT] 21 Aug 2016

Higher-order identities for balancing numbers

Takao Komatsu
School of Mathematics and Statistics
Wuhan University
Wuhan 430072 China
komatsu@whu.edu.cn

Prasanta Kumar Ray
VSS University of Technology, Odisha
Burla-768018, India
rayprasanta2008@gmail.com

Abstract

Let B_n be the n-th balancing number. In this paper, we give some explicit expressions of $\sum_{l=0}^{2r-3}(-1)^l\binom{2r-3}{l}\sum_{\substack{j_1+\cdots+j_r=n-2l\\ j_1,\ldots,j_r\geq 1}} B_{j_1}\cdots B_{j_r}$ and $\sum_{\substack{j_1+\cdots+j_r=n\\ j_1,\ldots,j_r\geq 1}} B_{j_1}\cdots B_{j_r}$. We also consider the convolution identities with binomial coefficients:

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 1}} \binom{n}{k_1,\ldots,k_r} B_{k_1}\cdots B_{k_r}$$

This type can be generalized, so that B_n is a special case of the number u_n, where $u_n = au_{n-1}+bu_{n-2}$ $(n\geq 2)$ with $u_0=0$ and $u_1=1$.

1 Introduction

Higher-order convolutions for various types of numbers (or polynomials) have been studied, with or without binomial (or multinomial) coefficients, including Bernoulli, Euler, Genocchi, Cauchy, Stirling, and Fibonacci numbers

([1, 2, 3, 6, 7, 8, 10]). One typical one is due to Euler, given by

$$\sum_{k=0}^{n} \binom{n}{k} \mathcal{B}_k \mathcal{B}_{n-k} = -n\mathcal{B}_{n-1} - (n-1)\mathcal{B}_n \quad (n \geq 0)\,,$$

where $\mathcal{B}_n$ are Bernoulli numbers, defined by

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} \mathcal{B}_n \frac{t^n}{n!} \quad (|t| < 2\pi)\,.$$

A positive integer x is called *balancing number* if

$$1 + 2 + \cdots + (x-1) = (x+1) + \cdots + (y-1) \tag{1}$$

holds for some integer $y \geq x + 2$. The problem of determining all balancing numbers leads to a Pell equation, whose solutions in x can be described by the recurrence $B_n = 6B_{n-1} - B_{n-2}$ $(n \geq 2)$ with $B_0 = 0$ and $B_1 = 1$ (see [4, 5]). One of the most general extensions of balancing numbers is when (1) is being replaced by

$$1^k + 2^k + \cdots + (x-1)^k = (x+1)^l + \cdots + (y-1)^l\,, \tag{2}$$

where the exponents k and l are given positive integers. In the work of Liptai et al. [12] effective and non-effective finiteness theorems on (2) are proved. In [9] a balancing problem of ordinary binomial coefficients is studied. Some more results can be seen in [13, 14, 15].

The generating function $f(x)$ of balancing numbers B_n is given by

$$f(x) := \frac{x}{1 - 6x + x^2} = \sum_{n=0}^{\infty} B_n x^n\,.$$

Then $f(x)$ satisfies the relation:

$$f(x)^2 = \frac{x^2}{1 - x^2} f'(x) \tag{3}$$

or

$$(1 - x^2) f(x)^2 = x^2 f'(x)\,. \tag{4}$$

The left-hand side of (4) is

$$\begin{aligned}
&(1-x^2)\left(\sum_{u=0}^{\infty} B_u x^u\right)\left(\sum_{v=0}^{\infty} B_v x^v\right) \\
&= (1-x^2)\sum_{n=0}^{\infty}\sum_{j=0}^{n} B_j B_{n-j} x^n \\
&= \sum_{n=0}^{\infty}\sum_{j=0}^{n} B_j B_{n-j} x^n - \sum_{n=2}^{\infty}\sum_{j=0}^{n-2} B_j B_{n-j-2} x^n \,.
\end{aligned}$$

The right-hand side of (4) is

$$x^2 \sum_{n=1}^{\infty} n B_n x^{n-1} = \sum_{n=0}^{\infty} (n-1) B_{n-1} x^n \,.$$

Comparing the coefficients of both sides, we get

$$\begin{aligned}
(n-1)B_{n-1} &= \sum_{j=0}^{n} B_j B_{n-j} - \sum_{j=0}^{n-2} B_j B_{n-j-2} \\
&= \sum_{j=1}^{n-1} (B_j B_{n-j} - B_{j-1} B_{n-j-1}) \,.
\end{aligned}$$

Here, notice that $B_0 = 0$. By changing n by $n+1$, we get the following identity.

Theorem 1. *For $n \geq 1$, we have*

$$n B_n = \sum_{j=1}^{n} (B_j B_{n-j+1} - B_{j-1} B_{n-j}) \,.$$

Differentiating both sides of (3) by x and dividing them by 2, we obtain

$$f(x) f'(x) = \frac{x}{(1-x^2)^2} f'(x) + \frac{x^2}{2(1-x^2)} f''(x) \,. \tag{5}$$

By (3) and (5), we get

$$\begin{aligned} f(x)^3 &= \frac{x^2}{1-x^2} f(x) f'(x) \\ &= \frac{x^3}{(1-x^2)^3} f'(x) + \frac{x^4}{2(1-x^2)^2} f''(x) \end{aligned} \tag{6}$$

or

$$(1-x^2)^3 f(x)^3 = x^3 f'(x) + \frac{1}{2} x^4 (1-x^2) f''(x)\,. \tag{7}$$

The left-hand side of (7) is equal to equal to

$$\begin{aligned} &(1 - 3x^2 + 3x^4 - x^6) \sum_{n=0}^{\infty} \sum_{\substack{j_1+j_2+J_3=n \\ j_1,j_2,j_3 \geq 0}} B_{j_1} B_{j_2} B_{j_3} x^n \\ &= \sum_{l=0}^{3} \sum_{n=2l}^{\infty} (-1)^l \binom{3}{l} \sum_{\substack{j_1+j_2+J_3=n-2l \\ j_1,j_2,j_3 \geq 1}} B_{j_1} B_{j_2} B_{j_3} x^n\,. \end{aligned}$$

The right-hand side of (7) is

$$\begin{aligned} &x^3 \sum_{n=1}^{\infty} n B_n x^{n-1} + \frac{x^4}{2} \sum_{n=2}^{\infty} n(n-1) B_n x^{n-2} - \frac{x^6}{2} \sum_{n=2}^{\infty} n(n-1) B_n x^{n-2} \\ &= \sum_{n=2}^{\infty} \frac{(n-1)(n-2)}{2} B_{n-2} x^n - \sum_{n=4}^{\infty} \frac{(n-4)(n-5)}{2} B_{n-4} x^n\,. \end{aligned}$$

Comparing the coefficients of both sides, we get the following result.

Theorem 2. *For $n \geq 4$, we have*

$$\sum_{l=0}^{3} (-1)^l \binom{3}{l} \sum_{\substack{j_1+j_2+J_3=n-2l \\ j_1,j_2,j_3 \geq 1}} B_{j_1} B_{j_2} B_{j_3} = \binom{n-1}{2} B_{n-2} - \binom{n-4}{2} B_{n-4}\,.$$

In this paper, we give some explicit expression of a more general case $\sum_{l=0}^{2r-3} (-1)^l \binom{2r-3}{l} \sum_{\substack{j_1+\cdots+j_r=n-2l \\ j_1,\ldots,j_r \geq 1}} B_{j_1} \cdots B_{j_r}$ and $\sum_{\substack{j_1+\cdots+j_r=n \\ j_1,\ldots,j_r \geq 1}} B_{j_1} \cdots B_{j_r}$. We also consider a different type for more general numbers:

$$\sum_{\substack{k_1+\cdots+k_r=n \\ k_1,\ldots,k_r \geq 1}} \binom{n}{k_1,\ldots,k_r} u_{k_1} \cdots u_{k_r}\,,$$

where $u_n = au_{n-1} + bu_{n-2}$ $(n \geq 2)$ with $u_0 = 0$ and $u_1 = 1$. When $a = 6$ and $b = -1$, this is reduced to the convolution identity for balancing numbers. When $a = b = 1$, this is reduced to the convolution identity for Fibonacci numbers. The corresponding identities for Luca-balancing numbers are also given.

2 Main results

First, as a general case of (3) and (6), we can have the following.

Lemma 1. *For $r \geq 2$, we have*

$$f(x)^r = \frac{x^{2r-2} f^{(r-1)}(x)}{(r-1)!(1-x^2)^{r-1}} + \sum_{k=1}^{r-2} \frac{\sum_{j=0}^{k-1} \binom{k}{j} \binom{r-2}{k-j-1} x^{2r-k+2j-2}}{k(r-k-2)!(1-x^2)^{r+k-1}} f^{(r-k-1)}(x)\,. \tag{8}$$

Proof. The proof is done by induction. It is trivial to see that the identity holds for $r = 2$. Suppose that the identity holds for some r. Differentiating

both sides by x, we obtain

$$
\begin{aligned}
&rf(x)^{r-1}f'(x)\\
&= \frac{x^{2r-2}f^{(r)}(x)}{(r-1)!(1-x^2)^{r-1}} + \frac{(2r-2)x^{2r-3}f^{(r-1)}(x)}{(r-1)!(1-x^2)^{r}}\\
&\quad + \sum_{k=1}^{r-2} \frac{\sum_{j=0}^{k-1} \binom{k}{j}\binom{r-2}{k-j-1} x^{2r-k-2+2j}}{k(r-k-2)!(1-x^2)^{r+k-1}} f^{(r-k)}(x)\\
&\quad + \sum_{k=1}^{r-2} \frac{\sum_{j=0}^{k-1} (2r-k-2+2j)\binom{k}{j}\binom{r-2}{k-j-1} x^{2r-k-3+2j}}{k(r-k-2)!(1-x^2)^{r+k}} f^{(r-k-1)}(x)\\
&\quad + \sum_{k=1}^{r-2} \frac{\sum_{j=0}^{k-1} (3k-2j)\binom{k}{j}\binom{r-2}{k-j-1} x^{2r-k-1+2j}}{k(r-k-2)!(1-x^2)^{r+k}} f^{(r-k-1)}(x)\\
&= \frac{x^{2r-2}f^{(r)}(x)}{(r-1)!(1-x^2)^{r-1}} + \frac{2x^{2r-3}f^{(r-1)}(x)}{(r-2)!(1-x^2)^{r}}\\
&\quad + \sum_{k=1}^{r-2} \frac{\sum_{j=0}^{k-1} \binom{k}{j}\binom{r-2}{k-j-1} x^{2r-k-2+2j}}{k(r-k-2)!(1-x^2)^{r+k-1}} f^{(r-k)}(x)\\
&\quad + \sum_{k=2}^{r-1} \frac{\sum_{j=0}^{k-2} (2r-k-1+2j)\binom{k-1}{j}\binom{r-2}{k-j-2} x^{2r-k-2+2j}}{(k-1)(r-k-1)!(1-x^2)^{r+k-1}} f^{(r-k)}(x)\\
&\quad + \sum_{k=2}^{r-1} \frac{\sum_{j=1}^{k-1} (3k-2j-1)\binom{k-1}{j-1}\binom{r-2}{k-j-1} x^{2r-k-2+2j}}{(k-1)(r-k-1)!(1-x^2)^{r+k-1}} f^{(r-k)}(x)\\
&= \frac{x^{2r-2}f^{(r)}(x)}{(r-1)!(1-x^2)^{r-1}} + r\sum_{k=1}^{r-1} \frac{\sum_{j=0}^{k-1} \binom{k}{j}\binom{r-1}{k-j-1} x^{2r-k-2+2j}}{k(r-k-1)!(1-x^2)^{r+k-1}} f^{(r-k)}(x)\,.
\end{aligned}
$$

Here, we used the relations

$$
\frac{2}{(r-2)!} + \frac{1}{(r-3)!} = \frac{r}{(r-2)!} \quad (k=1)
$$

and

$$\begin{aligned}
&\frac{r-k-1}{k}\binom{k}{j}\binom{r-2}{k-j-1}+\frac{2r-k-1+2j}{k-1}\binom{k-1}{j}\binom{r-2}{k-j-2}\\
&\qquad+\frac{3k-2j-1}{k-1}\binom{k-1}{j-1}\binom{r-2}{k-j-1}\\
&=\frac{r}{k}\binom{k}{j}\binom{r-1}{k-j-1}\quad(k\geq 2)\,.
\end{aligned}$$

Together with (3), we get

$$\begin{aligned}
f(x)^{r+1}&=\frac{x^2}{1-x^2}f(x)^{r-1}f'(x)\\
&=\frac{x^2}{1-x^2}\left(\frac{x^{2r-2}f^{(r)}(x)}{r!(1-x^2)^{r-1}}+\sum_{k=1}^{r-1}\frac{\sum_{j=0}^{k-1}\binom{k}{j}\binom{r-1}{k-j-1}x^{2r-k-2+2j}}{k(r-k-1)!(1-x^2)^{r+k-1}}f^{(r-k)}(x)\right)\\
&=\frac{x^{2r}f^{(r)}(x)}{r!(1-x^2)^r}+\sum_{k=1}^{r-1}\frac{\sum_{j=0}^{k-1}\binom{k}{j}\binom{r-1}{k-j-1}x^{2r-k+2j}}{k(r-k-1)!(1-x^2)^{r+k}}f^{(r-k)}(x)\,.
\end{aligned}$$

□

In general, we can state the following.

Theorem 3. *Let $r\geq 2$. Then for $n\geq 3r-5$, we have*

$$\begin{aligned}
&\sum_{l=0}^{2r-3}(-1)^l\binom{2r-3}{l}\sum_{\substack{j_1+\cdots+j_r=n-2l\\ j_1,\ldots,j_r\geq 1}}B_{j_1}\cdots B_{j_r}\\
&=\sum_{k=1}^{r-1}(-1)^{k-1}\frac{n-2k-r+3}{r-1}\binom{n-2k+1}{r-k-1}\binom{n-k-2r+3}{k-1}B_{n-2k-r+3}\,.
\end{aligned}$$

Proof. By Lemma 1 we get

$$\begin{aligned}
(1-x^2)^{2r-3}f(x)^r&=(1-x^2)^{r-2}\frac{x^{2r-2}f^{(r-1)}(x)}{(r-1)!}\\
&\quad+\sum_{k=1}^{r-2}(1-x^2)^{r-k-2}\frac{\sum_{j=0}^{k-1}\binom{k}{j}\binom{r-2}{k-j-1}x^{2r-k-2+2j}}{k(r-k-2)!}f^{(r-k-1)}(x)\,. \quad (9)
\end{aligned}$$

Since $B_0 = 0$, the left-hand side of (9) is equal to

$$\begin{aligned}
&(1-x^2)^{2r-3} \sum_{n=0}^{\infty} \sum_{\substack{j_1+\cdots+j_r=n \\ j_1,\ldots,j_r \geq 0}} B_{j_1} \cdots B_{j_r} x^n \\
&= \sum_{l=0}^{2r-3} \sum_{n=2l}^{\infty} (-1)^l \binom{2r-3}{l} \sum_{\substack{j_1+\cdots+j_r=n-2l \\ j_1,\ldots,j_r \geq 1}} B_{j_1} \cdots B_{j_r} x^n \,.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
&(1-x^2)^{r-2} \frac{x^{2r-2} f^{(r-1)}(x)}{(r-1)!} \\
&= \sum_{i=0}^{r-2} \binom{r-2}{i} x^{2i} \frac{x^{2r-2}}{(r-1)!} \sum_{n=r-1}^{\infty} \frac{n!}{(n-r+1)!} B_n x^{n-r+1} \\
&= \frac{1}{(r-1)!} \sum_{i=0}^{r-2} \binom{r-2}{i} \sum_{n=2r+2i-2}^{\infty} \frac{(n-r-2i+1)!}{(n-2r-2i+2)!} B_{n-r-2i+1} x^n \,.
\end{aligned}$$

For $i = r-2$, we have

$$\begin{aligned}
&\frac{(-1)^{r-2}}{(r-1)!} \sum_{n=4r-6}^{\infty} \frac{(n-3r+5)!}{(n-4r+6)!} B_{n-3r+5} x^n \\
&= (-1)^{r-2} \sum_{n=3r-5}^{\infty} \frac{n-3r+5}{r-1} \binom{n-3r+4}{r-2} B_{n-3r+5} x^n \,,
\end{aligned}$$

which yields the term for $k = r-1$ on the right-hand side of the identity in Theorem 3. Notice that

$$\binom{\gamma'}{\gamma} = 0 \quad (\gamma' < \gamma) \,.$$

The second term of the right-hand side of (9) is

$$\begin{aligned}
&\sum_{k=1}^{r-2}\sum_{i=0}^{r-k-2}(-1)^i\binom{r-k-2}{i}x^{2i}\frac{1}{k(r-k-2)!}\sum_{j=0}^{k-1}\binom{k}{j}\binom{r-2}{k-j-1}x^{2r-k-2+2j}\\
&\qquad\times\sum_{n=r-k-1}^{\infty}\frac{n!}{(n-r+k+1)!}B_n x^{n-r+k+1}\\
&=\sum_{i=0}^{r-3}\sum_{j=0}^{r-i-3}\sum_{k=j}^{r-i-3}\frac{(-1)^i}{(k+1)(r-k-3)!(n-2r+k-2i-2j+3)!}\binom{r-k-3}{i}\\
&\quad\times\binom{k+1}{j}\binom{r-2}{k-j}\sum_{n=2r+2i+2j-k-3}(n-r-2i-2j+1)!B_{n-r-2i-2j+1}x^n\\
&=\sum_{i=0}^{r-3}\sum_{\kappa=i+1}^{r-2}\sum_{k=\kappa-i-1}^{r-i-3}\frac{(-1)^i}{(k+1)(r-k-3)!(n-2r+k-2\kappa+5)!}\binom{r-k-3}{i}\\
&\quad\times\binom{k+1}{\kappa-i-1}\binom{r-2}{k-\kappa+i+1}\sum_{n=2r+2\kappa-k-5}(n-r-2\kappa+3)!B_{n-r-2\kappa+3}x^n\,.
\end{aligned}$$

Together with the first term of the right-hand side of (9) we can prove that

$$\begin{aligned}
&\frac{(-1)^{k-1}}{(r-1)!}\binom{r-2}{k-1}\frac{(n-r-2k+3)!}{(n-2r-2k+4)!}\\
&\quad+\sum_{i=0}^{k-1}\sum_{l=k-i-1}^{r-i-3}\frac{(-1)^{k-1}}{(l+1)(r-l-3)!(n-2r+l-2k+5)!}\\
&\qquad\times\binom{r-l-3}{i}\binom{l+1}{k-i-1}\binom{r-2}{l-k+i+1}(n-r-2k+3)!\\
&=(-1)^{k-1}\frac{n-2k-r+3}{r-1}\binom{n-2k+1}{r-k-1}\binom{n-k-2r+3}{k-1}. \qquad (10)
\end{aligned}$$

Then the proof is done. □

3 Examples

When $r=2$ and $r=3$, Theorem 3 is reduced to Theorem 1 and Theorem 2, respectively. When $r=4,5,6$ in Theorem 3, we get the following Corollaries as examples.

Corollary 1. *For $n \geq 7$, we have*

$$\sum_{l=0}^{5}(-1)^l\binom{5}{l}\sum_{\substack{j_1+j_2+j_3+j_4=n-2l\\ j_1,j_2,j_3,j_4\geq 1}} B_{j_1}B_{j_2}B_{j_3}B_{j_4}$$
$$= \binom{n-1}{3}B_{n-3} - \frac{(n-3)(n-5)(n-7)}{3}B_{n-5} + \binom{n-7}{3}B_{n-7}\,.$$

Corollary 2. *For $n \geq 10$, we have*

$$\sum_{l=0}^{7}(-1)^l\binom{7}{l}\sum_{\substack{j_1+j_2+j_3+j_4+j_5=n-2l\\ j_1,j_2,j_3,j_4,j_5\geq 1}} B_{j_1}B_{j_2}B_{j_3}B_{j_4}B_{j_5}$$
$$= \binom{n-1}{4}B_{n-4} - \frac{(n-3)(n-4)(n-6)(n-9)}{8}B_{n-6}$$
$$+ \frac{(n-5)(n-8)(n-10)(n-11)}{8}B_{n-6} - \binom{n-10}{4}B_{n-10}\,.$$

Corollary 3. *For $n \geq 13$, we have*

$$\sum_{l=0}^{9}(-1)^l\binom{9}{l}\sum_{\substack{j_1+\cdots+j_6=n-2l\\ j_1,\ldots,j_6\geq 1}} B_{j_1}\cdots B_{j_6}$$
$$= \binom{n-1}{5}B_{n-5} - \frac{(n-3)(n-4)(n-5)(n-7)(n-11)}{30}B_{n-7}$$
$$+ \frac{(n-5)(n-6)(n-9)(n-12)(n-13)}{20}B_{n-9}$$
$$- \frac{(n-7)(n-11)(n-13)(n-14)(n-15)}{30}B_{n-11} + \binom{n-13}{5}B_{n-13}\,.$$

4 Another result

In this section, we shall give an expression of $\sum_{\substack{j_1+\cdots+j_r=n\\ j_1,\ldots,j_r\geq 1}} B_{j_1}\cdots B_{j_r}$.

The left-hand side of (3) is equal to

$$\left(\sum_{u=0}^{\infty} B_u x^u\right)\left(\sum_{v=0}^{\infty} B_v x^v\right) = \sum_{n=0}^{\infty}\sum_{j=0}^{n} B_j B_{n-j} x^n \,.$$

The right-hand side of (3) is equal to

$$\begin{aligned}
&x^2\left(\sum_{j=0}^{\infty} x^{2j}\right)\left(\sum_{m=1}^{\infty} mB_m x^{m-1}\right)\\
&= x\left(\sum_{j=0}^{\infty} \frac{1+(-1)^j}{2} x^j\right)\left(\sum_{m=1}^{\infty} mB_m x^m\right)\\
&= x\sum_{n=0}^{\infty}\left(\sum_{m=0}^{n} \frac{1+(-1)^m}{2}(n-m)B_{n-m}\right)x^n\\
&= \sum_{n=1}^{\infty}\left(\sum_{m=0}^{n-1} \frac{1+(-1)^m}{2}(n-m-1)B_{n-m-1}\right)x^n\\
&= \sum_{n=1}^{\infty}\left(\sum_{m=0}^{\lfloor\frac{n-1}{2}\rfloor} (n-2m-1)B_{n-2m-1}\right)x^n \,.
\end{aligned}$$

Comparing the coefficients of both sides, we have the following.

Theorem 4. *For $n \geq 2$, we have*

$$\sum_{j=1}^{n-1} B_j B_{n-j} = \sum_{m=0}^{\lfloor\frac{n-1}{2}\rfloor} (n-2m-1)B_{n-2m-1} \,.$$

In general, we have the following.

Theorem 5. *For $n \geq r \geq 2$, we have*

$$\begin{aligned}
&\sum_{\substack{j_1+\cdots+j_r=n\\ j_1,\ldots,j_r\geq 1}} B_{j_1}\cdots B_{j_r}\\
&\qquad = \sum_{m=0}^{\lfloor\frac{n-r+1}{2}\rfloor} \binom{n-m-1}{r-2}\binom{m+r-2}{r-2}\frac{n-2m-r+1}{r-1}B_{n-2m-r+1} \,.
\end{aligned}$$

Proof. The left-hand side of (8) in Lemma 1 is equal to

$$\sum_{n=0}^{\infty} \sum_{\substack{j_1+\cdots+j_r=n \\ j_1,\ldots,j_r\geq 1}} B_{j_1}\cdots B_{j_r} x^n \,.$$

The first term on the right-hand side of (8) in Lemma 1 is equal to

$$\begin{aligned}
&\frac{x^{2r-2} f^{(r-1)}(x)}{(r-1)!(1-x^2)^{r-1}} \\
&= \frac{x^{2r-2}}{(r-1)!} \sum_{i=0}^{\infty} \binom{i+r-2}{r-2} x^{2i} \sum_{m=0}^{\infty} \frac{(m+r-1)!}{m!} B_{m+r-1} x^m \\
&= \frac{x^{2r-2}}{(r-1)!} \sum_{k=0}^{\infty} \frac{1}{(r-2)!2^{r-2}} \frac{(k+2r-4)!!}{k!!} \frac{1+(-1)^k}{2} x^k \\
&\quad \times \sum_{m=0}^{\infty} \frac{(m+r-1)!}{m!} B_{m+r-1} x^m \\
&= \frac{x^{2r-2}}{(r-1)!} \sum_{n=0}^{\infty} \sum_{m=0}^{n} \frac{1}{(r-2)!2^{r-2}} \frac{(n-m+2r-4)!!}{(n-m)!!} \\
&\qquad \times \frac{1+(-1)^{n-m}}{2} \frac{(m+r-1)!}{m!} B_{m+r-1} x^n \\
&= \frac{1}{(r-1)!(r-2)!2^{r-2}} \sum_{n=2r-2}^{\infty} \sum_{m=0}^{n-2r+2} \frac{(n-m-2)!!}{(n-m-2r+2)!!} \\
&\qquad \times \frac{1+(-1)^{n-m}}{2} \frac{(m+r-1)!}{m!} B_{m+r-1} x^n \\
&= \frac{1}{(r-1)!(r-2)!2^{r-2}} \sum_{n=2r-2}^{\infty} \sum_{m=r-1}^{n-r+1} \frac{(n-m+r-3)!!}{(n-m-r+1)!!} \\
&\qquad \times \frac{1+(-1)^{n-m-r+1}}{2} \frac{m!}{(m-r+1)!} B_m x^n \,.
\end{aligned}$$

Concerning the second term, we have

$$
\begin{aligned}
&\frac{\sum_{j=0}^{k-1}\binom{k}{j}\binom{r-2}{k-j-1}x^{2r-k-2+2j}}{(1-x^2)^{r+k-1}}f^{(r-k-1)}(x)\\
&=\sum_{j=0}^{k-1}\binom{k}{j}\binom{r-2}{k-j-1}x^{2r-k-2+2j}\sum_{i=0}^{\infty}\binom{i+r+k-2}{r+k-2}x^{2i}\\
&\qquad\times\sum_{m=0}^{\infty}\frac{(m+r-k-1)!}{m!}B_{m+r-k-1}x^m\\
&=\sum_{j=0}^{k-1}\binom{k}{j}\binom{r-2}{k-j-1}x^{2r-k-2+2j}\sum_{l=0}^{\infty}\frac{1}{(r+k-2)!2^{r+k-2}}\\
&\qquad\times\frac{(l+2r+2k-4)!!}{l!!}\frac{1+(-1)^l}{2}x^l\sum_{m=0}^{\infty}\frac{(m+r-k-1)!}{m!}B_{m+r-k-1}x^m\\
&=\sum_{j=0}^{k-1}\binom{k}{j}\binom{r-2}{k-j-1}x^{2r-k-2+2j}\sum_{n=0}^{\infty}\sum_{m=0}^{n}\frac{1}{(r+k-2)!2^{r+k-2}}\\
&\qquad\times\frac{(n-m+2r+2k-4)!!}{(n-m)!!}\frac{1+(-1)^{n-m}}{2}\frac{(m+r-k-1)!}{m!}B_{m+r-k-1}x^n\\
&=\frac{1}{(r+k-2)!2^{r+k-2}}\sum_{j=0}^{k-1}\binom{k}{j}\binom{r-2}{k-j-1}\sum_{n=2r-k-2+2j}^{\infty}\sum_{m=0}^{n-2r+k+2-2j}\\
&\quad\frac{(n-m+3k-2-2j)!!}{(n-m-2r+k+2-2j)!!}\frac{1+(-1)^{n-m+k}}{2}\frac{(m+r-k-1)!}{m!}B_{m+r-k-1}x^n\,.
\end{aligned}
$$

Since

$$\frac{(n-m+r+2k-3-2j)!!}{(n-m-r+k+3-2j)!!}=0$$

if $m = n - 2r + k + 2 - 2j$ $(j = 1, 2, \ldots, k-2)$, this is equal to

$$\begin{aligned}
&\frac{1}{(r+k-2)!2^{r+k-2}} \sum_{j=0}^{k-1} \binom{k}{j} \binom{r-2}{k-j-1} \sum_{n=2r-k-2}^{\infty} \sum_{m=0}^{n-2r+k+2} \\
&\times \frac{(n-m+3k-2-2j)!!}{(n-m+k)!!} \frac{1+(-1)^{n-m+k}}{2} \frac{(m+r-k-1)!}{m!} B_{m+r-k-1} x^n \\
&= \frac{1}{(r+k-2)!2^{r+k-2}} \sum_{n=2r-k-2}^{\infty} \sum_{m=r-k-1}^{n-r+1} \sum_{j=0}^{k-1} \binom{k}{j} \binom{r-2}{k-j-1} \\
&\quad \times \frac{(n-m+r+2k-3-2j)!!}{(n-m-r+k+3-2j)!!} \frac{1+(-1)^{n-m-r+1}}{2} \frac{m!}{(m-r+k+1)!} B_m x^n \,.
\end{aligned}$$

This is also equal to

$$\begin{aligned}
&\frac{1}{(r+k-2)!2^{r+k-2}} \sum_{n=2r-k-2}^{\infty} \sum_{m=r-k-1}^{n-r+1} \sum_{j=0}^{k-1} \binom{k}{j} \binom{r-2}{k-j-1} \\
&\quad \times \frac{(n-m+r+2k-3-2j)!!}{(n-m-r+k+3-2j)!!} \frac{1+(-1)^{n-m-r+1}}{2} \frac{m!}{(m-r+k+1)!} B_m x^n \\
&= \frac{1}{(r+k-2)!2^{r+k-2}} \sum_{n=2r-k-2}^{\infty} \sum_{m=r-k-1}^{n-r+1} \frac{(n-m+r-1)!!}{(n-m-r+1)!!} \binom{r+k-2}{k-1} \\
&\quad \times \frac{(n-m+r-3)!!}{(n-m+r-2k-1)!!} \frac{1+(-1)^{n-m-r+1}}{2} \frac{m!}{(m-r+k+1)!} B_m x^n \,.
\end{aligned}$$

Therefore, the right-hand side of the relation in Theorem 4 is

$$\begin{aligned}
&\frac{1}{(r-1)!(r-2)!2^{r-2}}\sum_{n=2r-2}^{\infty}\sum_{m=r-1}^{n-r+1}\frac{(n-m+r-3)!!}{(n-m-r+1)!!}\frac{1+(-1)^{n-m-r+1}}{2}\\
&\qquad\times\frac{m!}{(m-r+1)!}B_m x^n\\
&\quad+\sum_{k=1}^{r-1}\frac{1}{k(r-k-2)!}\frac{1}{(r+k-2)!2^{r+k-2}}\sum_{n=2r-k-2}^{\infty}\\
&\qquad\sum_{m=r-k-1}^{n-r+1}\frac{(n-m+r-1)!!}{(n-m-r+1)!!}\binom{r+k-2}{k-1}\frac{(n-m+r-3)!!}{(n-m+r-2k-1)!!}\\
&\qquad\qquad\times\frac{1+(-1)^{n-m-r+1}}{2}\frac{m!}{(m-r+k+1)!}B_m x^n\\
&=\frac{1}{(r-1)!(r-2)!2^{r-2}}\sum_{n=2r-2}^{\infty}\sum_{m=r-1}^{n-r+1}\frac{(n-m+r-3)!!}{(n-m-r+1)!!}\frac{1+(-1)^{n-m-r+1}}{2}\\
&\qquad\times\frac{m!}{(m-r+1)!}B_m x^n\\
&\quad+\sum_{n=r-1}^{\infty}\frac{1}{(r-1)!2^{r-2}}\sum_{m=1}^{r-2}\sum_{k=r-m-1}^{r-2}\frac{1}{k!(r-k-2)!2^k}\frac{(n-m+r-1)!!}{(n-m-r+1)!!}\\
&\qquad\times\frac{(n-m+r-3)!!}{(n-m+r-2k-1)!!}\frac{1+(-1)^{n-m-r+1}}{2}\frac{m!}{(m-r+k+1)!}B_m x^n\\
&\quad+\sum_{n=r-1}^{\infty}\frac{1}{(r-1)!2^{r-2}}\sum_{m=r-1}^{n-r+1}\sum_{k=1}^{r-2}\frac{1}{k!(r-k-2)!2^k}\frac{(n-m+r-1)!!}{(n-m-r+1)!!}\\
&\qquad\times\frac{(n-m+r-3)!!}{(n-m+r-2k-1)!!}\frac{1+(-1)^{n-m-r+1}}{2}\frac{m!}{(m-r+k+1)!}B_m x^n\,.
\end{aligned}$$

Since for $1 \le m \le r-2$ we have

$$\begin{aligned}
&\frac{1}{(r-1)!2^{r-2}} \sum_{k=r-m-1}^{r-2} \frac{1}{k!(r-k-2)!2^k} \frac{(n-m+r-1)!!}{(n-m-r+1)!!} \\
&\qquad \times \frac{(n-m+r-3)!!}{(n-m+r-2k-1)!!} \frac{m!}{(m-r+k+1)!} \\
&= \frac{1}{(r-1)!(r-2)!2^{2r-4}} \frac{(n+m+r-3)!!}{(n+m-r+1)!!} \frac{(n-m+r-3)!!}{(n-m-r+1)!!} m
\end{aligned}$$

and for $r-1 \le m \le n-r+1$ we have

$$\begin{aligned}
&\frac{1}{(r-1)!(r-2)!2^{r-2}} \frac{(n-m+r-3)!!}{(n-m-r+1)!!} \frac{m!}{(m-r+1)!} \\
&\quad + \frac{1}{(r-1)!2^{r-2}} \sum_{k=r-m-1}^{r-2} \frac{1}{k!(r-k-2)!2^k} \frac{(n-m+r-1)!!}{(n-m-r+1)!!} \\
&\qquad \times \frac{(n-m+r-3)!!}{(n-m+r-2k-1)!!} \frac{m!}{(m-r+k+1)!} \\
&= \frac{1}{(r-1)!(r-2)!2^{2r-4}} \frac{(n+m+r-3)!!}{(n+m-r+1)!!} \frac{(n-m+r-3)!!}{(n-m-r+1)!!} m \,,
\end{aligned}$$

By comparing the coefficients, we have

$$\begin{aligned}
&\sum_{\substack{j_1+\cdots+j_r=n\\ j_1,\ldots,j_r\geq 1}} B_{j_1}\cdots B_{j_r}\\
&= \frac{1}{(r-1)!(r-2)!2^{2r-4}}\\
&\quad\times \sum_{m=0}^{n-r+1} \frac{(n+m+r-3)!!(n-m+r-3)!!}{(n+m-r+1)!!(n-m-r+1)!!}\frac{1+(-1)^{n-m-r+1}}{2}mF_m\\
&= \frac{1}{(r-1)!(r-2)!2^{2r-4}}\\
&\quad\times \sum_{m=0}^{n-r+1} \frac{(2n-m-2)!!(m+2r-4)!!}{(2n-m-2r+2)!!m!!}(n-m-r+1)B_{n-m-r+1}\\
&= \frac{1}{(r-1)!(r-2)!2^{2r-4}}\\
&\quad\times \sum_{m=0}^{\lfloor\frac{n-r+1}{2}\rfloor} \frac{(2n-2m-2)!!(2m+2r-4)!!}{(2n-2m-2r+2)!!(2m)!!}(n-2m-r+1)B_{n-2m-r+1}\\
&= \sum_{m=0}^{\lfloor\frac{n-r+1}{2}\rfloor} \binom{n-m-1}{r-2}\binom{m+r-2}{r-2}\frac{n-2m-r+1}{r-1}B_{n-2m-r+1}\,.
\end{aligned}$$

□

5 Some other generating functions

Another kinds of the generating functions of balancing numbers and Lucas-balancing numbers are given by

$$b(t) := \frac{e^{\alpha t}-e^{\beta t}}{4\sqrt{2}} = \sum_{n=0}^{\infty} B_n\frac{t^n}{n!}$$

and

$$c(t) := \frac{e^{\alpha t}+e^{\beta t}}{2} = \sum_{n=0}^{\infty} C_n\frac{t^n}{n!}$$

because they satisfy the differential equation $y''-6y'+y=0$.

Since $b'(t) = 3b(t) + c(t)$ and $c'(t) = 8b(t) + 3c(t)$, we have for $n \geq 0$

$$B_{n+1} = 3B_n + C_n$$

and

$$C_{n+1} = 8B_n + 3C_n\,.$$

Since

$$c(t)^2 = \frac{e^{2\alpha t} + e^{2\beta t}}{4} + \frac{e^{6t}}{2}\,,$$

we have

$$\sum_{n=0}^{\infty}\sum_{k=0}^{n}\binom{n}{k}C_kC_{n-k} = \frac{1}{2}\sum_{n=0}^{\infty}C_n\frac{(2t)^n}{n!} + \frac{1}{2}\sum_{n=0}^{\infty}\frac{(6t)^n}{n!}\,,$$

yielding

$$\sum_{k=0}^{n}\binom{n}{k}C_kC_{n-k} = \frac{2^nC_n + 6^n}{2} \quad (n \geq 0)\,.$$

Similarly, by

$$b(t)^2 = \frac{e^{2\alpha t} + e^{2\beta t}}{32} - \frac{e^{6t}}{16}\,,$$

we have

$$\sum_{k=0}^{n}\binom{n}{k}B_kB_{n-k} = \frac{2^nC_n - 6^n}{16} \quad (n \geq 0)\,.$$

Since by $2\alpha + \beta = \alpha + 6$ and $\alpha + 2\beta = \beta + 6$

$$\begin{aligned}
b(t)^3 &= \frac{e^{3\alpha t} - e^{3\beta t}}{(4\sqrt{2})^3} - \frac{3(e^{(2\alpha+\beta)t} - e^{(\alpha+2\beta)t})}{(4\sqrt{2})^3} \\
&= \frac{1}{32}\frac{e^{3\alpha t} - e^{3\beta t}}{4\sqrt{2}} - \frac{3}{32}e^{6t}\frac{e^{\alpha t} - e^{\beta t}}{4\sqrt{2}} \\
&= \frac{1}{32}\sum_{n=0}^{\infty}B_n\frac{(3t)^n}{n!} - \frac{3}{32}\sum_{i=0}^{\infty}\frac{(6t)^i}{i!}\sum_{k=0}^{\infty}B_k\frac{t^k}{k!} \\
&= \frac{1}{32}\sum_{n=0}^{\infty}3^nB_n\frac{t^n}{n!} - \frac{3}{32}\sum_{n=0}^{\infty}\sum_{k=0}^{n}\binom{n}{k}6^{n-k}B_k\frac{t^n}{n!}\,,
\end{aligned}$$

we have

$$\sum_{\substack{k_1+k_2+k_3=n\\ k_1,k_2,k_3\geq 1}} \binom{n}{k_1,k_2,k_3} B_{k_1}B_{k_2}B_{k_3} = \frac{1}{32}\left(3^n B_n - 3\sum_{k=0}^{n}\binom{n}{k}6^{n-k}B_k\right).$$

Notice that $B_0 = 0$. Similarly, we can obtain that

$$\sum_{\substack{k_1+k_2+k_3=n\\ k_1,k_2,k_3\geq 0}} \binom{n}{k_1,k_2,k_3} C_{k_1}C_{k_2}C_{k_3} = \frac{1}{4}\left(3^n C_n + 3\sum_{k=0}^{n}\binom{n}{k}6^{n-k}C_k\right).$$

Let $r \geq 1$. If r is odd, then

$$\begin{aligned}
b(t)^r &= \left(\frac{e^{\alpha t} - e^{\beta t}}{4\sqrt{2}}\right)^r \\
&= \frac{1}{(4\sqrt{2})^r}\sum_{j=0}^{\frac{r-1}{2}}(-1)^j\binom{r}{j}(e^{((r-j)\alpha+j\beta)t} - e^{(j\alpha+(r-j)\beta)t}) \\
&= \frac{1}{(4\sqrt{2})^r}\sum_{j=0}^{\frac{r-1}{2}}(-1)^j\binom{r}{j}e^{6jt}(e^{(r-2j)\alpha t} - e^{(r-2j)\beta t}) \\
&= \frac{1}{(4\sqrt{2})^{r-1}}\sum_{j=0}^{\frac{r-1}{2}}(-1)^j\binom{r}{j}\sum_{i=0}^{\infty}\frac{(6jt)^i}{i!}\sum_{k=0}^{\infty}B_k\frac{\big((r-2j)t\big)^k}{k!} \\
&= \frac{1}{(4\sqrt{2})^{r-1}}\sum_{j=0}^{\frac{r-1}{2}}(-1)^j\binom{r}{j}\sum_{n=0}^{\infty}\sum_{k=0}^{n}\binom{n}{k}(6j)^{n-k}(r-2j)^k B_k\frac{t^n}{n!}.
\end{aligned}$$

Therefore, we get

$$\begin{aligned}
&\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 1}} \binom{n}{k_1,\ldots,k_r} B_{k_1}\cdots B_{k_r} \\
&\qquad = \frac{1}{(4\sqrt{2})^{r-1}}\sum_{j=0}^{\frac{r-1}{2}}(-1)^j\binom{r}{j}\sum_{k=0}^{n}\binom{n}{k}(6j)^{n-k}(r-2j)^k B_k.
\end{aligned}$$

Similarly, we get

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 0}} \binom{n}{k_1,\ldots,k_r} C_{k_1}\cdots C_{k_r}$$

$$= \frac{1}{2^{r-1}} \sum_{j=0}^{\frac{r-1}{2}} \binom{r}{j} \sum_{k=0}^{n} \binom{n}{k} (6j)^{n-k} (r-2j)^k C_k \,.$$

If r is even, then

$$\begin{aligned}
b(t)^r &= \frac{1}{(4\sqrt{2})^r} \left(\sum_{j=0}^{\frac{r}{2}-1} (-1)^j \binom{r}{j} (e^{((r-j)\alpha+j\beta)t} + e^{(j\alpha+(r-j)\beta)t}) \right. \\
&\quad \left. +(-1)^{\frac{r}{2}} \binom{r}{\frac{r}{2}} e^{(\frac{r}{2}\alpha+\frac{r}{2}\beta)t} \right) \\
&= \frac{1}{(4\sqrt{2})^r} \left(\sum_{j=0}^{\frac{r}{2}-1} (-1)^j \binom{r}{j} e^{6jt} (e^{(r-2j)\alpha t} + e^{(r-2j)\beta t}) \right. \\
&\quad \left. +(-1)^{\frac{r}{2}} \binom{r}{\frac{r}{2}} e^{3rt} \right) \\
&= \frac{1}{(4\sqrt{2})^r} \left(\sum_{j=0}^{\frac{r}{2}-1} (-1)^j \binom{r}{j} \sum_{i=0}^{\infty} \frac{(6jt)^i}{i!} \cdot 2 \sum_{k=0}^{\infty} C_k \frac{\big((r-2j)t\big)^k}{k!} \right. \\
&\quad \left. +(-1)^{\frac{r}{2}} \binom{r}{\frac{r}{2}} \sum_{n=0}^{\infty} \frac{(3r)^n}{n!} t^n \right) \\
&= \frac{1}{(4\sqrt{2})^r} \left(2 \sum_{j=0}^{\frac{r}{2}-1} (-1)^j \binom{r}{j} \sum_{n=0}^{\infty} \sum_{k=0}^{n} \binom{n}{k} (6j)^{n-k} (r-2j)^k C_k \frac{t^n}{n!} \right. \\
&\quad \left. +(-1)^{\frac{r}{2}} \binom{r}{\frac{r}{2}} \sum_{n=0}^{\infty} (3r)^n \frac{t^n}{n!} \right) .
\end{aligned}$$

Therefore, we get

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 1}} \binom{n}{k_1,\ldots,k_r} B_{k_1}\cdots B_{k_r}$$
$$= \frac{1}{(4\sqrt{2})^r}\left(2\sum_{j=0}^{\frac{r}{2}-1}(-1)^j\binom{r}{j}\sum_{k=0}^{n}\binom{n}{k}(6j)^{n-k}(r-2j)^k C_k \right.$$
$$\left. +(-1)^{\frac{r}{2}}\binom{r}{\frac{r}{2}}(3r)^n\right).$$

Similarly, we get

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 0}} \binom{n}{k_1,\ldots,k_r} C_{k_1}\cdots C_{k_r}$$
$$= \frac{1}{2^r}\left(2\sum_{j=0}^{\frac{r}{2}-1}\binom{r}{j}\sum_{k=0}^{n}\binom{n}{k}(6j)^{n-k}(r-2j)^k C_k + \binom{r}{\frac{r}{2}}(3r)^n\right).$$

6 More general cases

Let $\{u_n\}_{n\geq 0}$ and $\{v_n\}_{n\geq 0}$ be integer sequences, satisfying the same recurrence relation: $u_n = au_{n-1} + bu_{n-2}$ $(n \geq 2)$ and $v_n = av_{n-1} + bv_{n-2}$ $(n \geq 2)$ with initial values u_0, u_1, v_0 and v_1. If the general terms are given by

$$u_n = \frac{\alpha^n - \beta^n}{\alpha - \beta} \quad \text{and} \quad v_n = \alpha^n + \beta^n \quad (n \geq 0)$$

where

$$\alpha = \frac{a + \sqrt{a^2+4b}}{2} \quad \text{and} \quad \beta = \frac{a - \sqrt{a^2+4b}}{2},$$

we can set $u_0 = 0$, $u_1 = 1$, $v = 0 = 2$ and $v_1 = a$. Then, the generating functions of u_n and v_n are given by

$$u(t) := \frac{e^{\alpha t} - e^{\beta t}}{\sqrt{a^2+4b}} = \sum_{n=0}^{\infty} u_n \frac{t^n}{n!} \quad \text{and} \quad v(t) := e^{\alpha t} + e^{\beta t} = \sum_{n=0}^{\infty} v_n \frac{t^n}{n!}$$

respectively, because they satisfy the differential equation $y'' - ay' - by = 0$.

Our main results can be stated as follows.

Theorem 6. *If r is odd, then*

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 1}} \binom{n}{k_1,\ldots,k_r} u_{k_1}\cdots u_{k_r}$$

$$= \frac{1}{(\sqrt{a^2+4b})^{r-1}} \sum_{k=0}^{n} \binom{n}{k} \sum_{j=0}^{\frac{r-1}{2}} (-1)^j \binom{r}{j} (aj)^{n-k}(r-2j)^k u_k \quad (11)$$

and

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 0}} \binom{n}{k_1,\ldots,k_r} v_{k_1}\cdots v_{k_r}$$

$$= \sum_{k=0}^{n} \binom{n}{k} \sum_{j=0}^{\frac{r-1}{2}} \binom{r}{j} (aj)^{n-k}(r-2j)^k v_k\,. \quad (12)$$

If r is even, then

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 1}} \binom{n}{k_1,\ldots,k_r} u_{k_1}\cdots u_{k_r}$$

$$= \frac{1}{(\sqrt{a^2+4b})^{r}} \left(\sum_{k=0}^{n} \binom{n}{k} \sum_{j=0}^{\frac{r}{2}-1} (-1)^j \binom{r}{j} (aj)^{n-k}(r-2j)^k v_k \right.$$

$$\left. +(-1)^{\frac{r}{2}} \binom{r}{\frac{r}{2}} \left(\frac{ar}{2}\right)^n \right) \quad (13)$$

and

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 0}} \binom{n}{k_1,\ldots,k_r} v_{k_1}\cdots v_{k_r}$$

$$= \sum_{k=0}^{n} \binom{n}{k} \sum_{j=0}^{\frac{r}{2}-1} \binom{r}{j} (aj)^{n-k}(r-2j)^k v_k + \binom{r}{\frac{r}{2}} \left(\frac{ar}{2}\right)^n\,. \quad (14)$$

Proof. If r is odd, then by $\alpha+\beta=a$

$$\begin{aligned}
u(t)^r &= \left(\frac{e^{\alpha t}-e^{\beta t}}{\sqrt{a^2+4b}}\right)^r \\
&= \frac{1}{(\sqrt{a^2+4b})^r}\sum_{j=0}^{\frac{r-1}{2}}(-1)^j\binom{r}{j}(e^{((r-j)\alpha+j\beta)t}-e^{(j\alpha+(r-j)\beta)t}) \\
&= \frac{1}{(\sqrt{a^2+4b})^r}\sum_{j=0}^{\frac{r-1}{2}}(-1)^j\binom{r}{j}e^{ajt}(e^{(r-2j)\alpha t}-e^{(r-2j)\beta t}) \\
&= \frac{1}{(\sqrt{a^2+4b})^{r-1}}\sum_{j=0}^{\frac{r-1}{2}}(-1)^j\binom{r}{j}\sum_{i=0}^{\infty}\frac{(ajt)^i}{i!}\sum_{k=0}^{\infty}u_k\frac{\big((r-2j)t\big)^k}{k!} \\
&= \frac{1}{(\sqrt{a^2+4b})^{r-1}}\sum_{j=0}^{\frac{r-1}{2}}(-1)^j\binom{r}{j}\sum_{n=0}^{\infty}\sum_{k=0}^{n}\binom{n}{k}(aj)^{n-k}(r-2j)^k u_k\frac{t^n}{n!}\,.
\end{aligned}$$

Therefore, we get

$$\begin{aligned}
&\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 1}}\binom{n}{k_1,\ldots,k_r}u_{k_1}\cdots u_{k_r} \\
&\qquad = \frac{1}{(\sqrt{a^2+4b})^{r-1}}\sum_{j=0}^{\frac{r-1}{2}}(-1)^j\binom{r}{j}\sum_{k=0}^{n}\binom{n}{k}(aj)^{n-k}(r-2j)^k u_k\,.
\end{aligned}$$

Similarly, we get

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 0}}\binom{n}{k_1,\ldots,k_r}v_{k_1}\cdots v_{k_r} = \sum_{j=0}^{\frac{r-1}{2}}\binom{r}{j}\sum_{k=0}^{n}\binom{n}{k}(aj)^{n-k}(r-2j)^k v_k\,.$$

If r is even, then

$$\begin{aligned}
u(t)^r &= \frac{1}{(\sqrt{a^2+4b})^r}\left(\sum_{j=0}^{\frac{r}{2}-1}(-1)^j\binom{r}{j}(e^{((r-j)\alpha+j\beta)t}+e^{(j\alpha+(r-j)\beta)t})\right.\\
&\quad\left.+(-1)^{\frac{r}{2}}\binom{r}{\frac{r}{2}}e^{(\frac{r}{2}\alpha+\frac{r}{2}\beta)t}\right)\\
&= \frac{1}{(\sqrt{a^2+4b})^r}\left(\sum_{j=0}^{\frac{r}{2}-1}(-1)^j\binom{r}{j}e^{ajt}(e^{(r-2j)\alpha t}+e^{(r-2j)\beta t})\right.\\
&\quad\left.+(-1)^{\frac{r}{2}}\binom{r}{\frac{r}{2}}e^{art/2}\right)\\
&= \frac{1}{(\sqrt{a^2+4b})^r}\left(\sum_{j=0}^{\frac{r}{2}-1}(-1)^j\binom{r}{j}\sum_{i=0}^{\infty}\frac{(ajt)^i}{i!}\cdot\sum_{k=0}^{\infty}v_k\frac{\big((r-2j)t\big)^k}{k!}\right.\\
&\quad\left.+(-1)^{\frac{r}{2}}\binom{r}{\frac{r}{2}}\sum_{n=0}^{\infty}\Big(\frac{ar}{2}\Big)^n\frac{t^n}{n!}\right)\\
&= \frac{1}{(\sqrt{a^2+4b})^r}\left(\sum_{j=0}^{\frac{r}{2}-1}(-1)^j\binom{r}{j}\sum_{n=0}^{\infty}\sum_{k=0}^{n}\binom{n}{k}(aj)^{n-k}(r-2j)^kv_k\frac{t^n}{n!}\right.\\
&\quad\left.+(-1)^{\frac{r}{2}}\binom{r}{\frac{r}{2}}\sum_{n=0}^{\infty}\Big(\frac{ar}{2}\Big)^n\frac{t^n}{n!}\right).
\end{aligned}$$

Therefore, we get

$$\begin{aligned}
&\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 1}}\binom{n}{k_1,\ldots,k_r}u_{k_1}\cdots u_{k_r}\\
&\qquad= \frac{1}{(\sqrt{a^2+4b})^r}\left(\sum_{j=0}^{\frac{r}{2}-1}(-1)^j\binom{r}{j}\sum_{k=0}^{n}\binom{n}{k}(aj)^{n-k}(r-2j)^kv_k\right.\\
&\qquad\qquad\left.+(-1)^{\frac{r}{2}}\binom{r}{\frac{r}{2}}\Big(\frac{ar}{2}\Big)^n\right).
\end{aligned}$$

Similarly, we get

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 0}} \binom{n}{k_1,\ldots,k_r} v_{k_1}\cdots v_{k_r}$$
$$= \sum_{j=0}^{\frac{r}{2}-1} \binom{r}{j} \sum_{k=0}^{n} \binom{n}{k} (aj)^{n-k}(r-2j)^k v_k + \binom{r}{\frac{r}{2}} \left(\frac{ar}{2}\right)^n .$$

□

7 Examples

When $a = 6$ and $b = -1$, then $B_n = u_n$ are balancing numbers and $C_n = v_n/2$ are Lucas-balancing numbers. Thus, Theorem 6 can be reduced as follows.

Corollary 4. *If r is odd, then*

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 1}} \binom{n}{k_1,\ldots,k_r} B_{k_1}\cdots B_{k_r}$$
$$= \frac{1}{(4\sqrt{2})^{r-1}} \sum_{k=0}^{n} \binom{n}{k} \sum_{j=0}^{\frac{r-1}{2}} (-1)^j \binom{r}{j} (6j)^{n-k}(r-2j)^k B_k$$

and

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 0}} \binom{n}{k_1,\ldots,k_r} C_{k_1}\cdots C_{k_r}$$
$$= \frac{1}{2^{r-1}} \sum_{k=0}^{n} \binom{n}{k} \sum_{j=0}^{\frac{r-1}{2}} \binom{r}{j} (6j)^{n-k}(r-2j)^k C_k .$$

If r is even, then

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 1}} \binom{n}{k_1,\ldots,k_r} B_{k_1}\cdots B_{k_r}$$
$$= \frac{1}{(4\sqrt{2})^r}\left(2\sum_{k=0}^{n}\binom{n}{k}\sum_{j=0}^{\frac{r}{2}-1}(-1)^j\binom{r}{j}(6j)^{n-k}(r-2j)^k C_k \right.$$
$$\left. +(-1)^{\frac{r}{2}}\binom{r}{\frac{r}{2}}\left(\frac{r}{2}\right)^n\right)$$

and

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 0}} \binom{n}{k_1,\ldots,k_r} C_{k_1}\cdots C_{k_r}$$
$$= \frac{1}{2^r}\left(2\sum_{k=0}^{n}\binom{n}{k}\sum_{j=0}^{\frac{r}{2}-1}\binom{r}{j}(6j)^{n-k}(r-2j)^k C_k + \binom{r}{\frac{r}{2}}\left(\frac{6r}{2}\right)^n\right).$$

When $a=b=1$, then $F_n=u_n$ are Fibonacci numbers and $L_n=v_n$ are Lucas numbers. Thus, Theorem 6 can be reduced as follows.

Corollary 5. *If r is odd, then*

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 1}} \binom{n}{k_1,\ldots,k_r} F_{k_1}\cdots F_{k_r}$$
$$= \frac{1}{(\sqrt{5})^{r-1}}\sum_{k=0}^{n}\binom{n}{k}\sum_{j=0}^{\frac{r-1}{2}}(-1)^j\binom{r}{j}j^{n-k}(r-2j)^k F_k$$

and

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 0}} \binom{n}{k_1,\ldots,k_r} L_{k_1}\cdots L_{k_r}$$
$$= \sum_{k=0}^{n}\binom{n}{k}\sum_{j=0}^{\frac{r-1}{2}}\binom{r}{j}j^{n-k}(r-2j)^k L_k\,.$$

If r is even, then

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 1}} \binom{n}{k_1,\ldots,k_r} F_{k_1}\cdots F_{k_r}$$
$$= \frac{1}{(\sqrt{5})^r}\left(\sum_{k=0}^{n}\binom{n}{k}\sum_{j=0}^{\frac{r}{2}-1}(-1)^j\binom{r}{j}j^{n-k}(r-2j)^k L_k \right.$$
$$\left. +(-1)^{\frac{r}{2}}\binom{r}{\frac{r}{2}}\left(\frac{r}{2}\right)^n\right)$$

and

$$\sum_{\substack{k_1+\cdots+k_r=n\\ k_1,\ldots,k_r\geq 0}} \binom{n}{k_1,\ldots,k_r} L_{k_1}\cdots L_{k_r} = \sum_{k=0}^{n}\binom{n}{k}\sum_{j=0}^{\frac{r}{2}-1}\binom{r}{j}j^{n-k}(r-2j)^k L_k + \binom{r}{\frac{r}{2}}\left(\frac{r}{2}\right)^n .$$

When $r=2$ and $a=b=1$, we have

$$\sum_{n=0}^{n}\binom{n}{k}F_k F_{n-k} = \frac{2^n L_n - 2}{5}$$

and

$$\sum_{n=0}^{n}\binom{n}{k}L_k L_{n-k} = 2^n L_n + 2\,.$$

References

[1] T. Agoh and K. Dilcher, *Convolution identities and lacunary recurrences for Bernoulli numbers*, J. Number Theory 124 (2007), 105–122.

[2] T. Agoh and K. Dilcher, *Higher-order recurrences for Bernoulli numbers*, J. Number Theory **129** (2009), 1837–1847.

[3] T. Agoh and K. Dilcher, *Higher-order convolutions for Bernoulli and Euler polynomials*, J. Math. Anal. Appl. **419** (2014), 1235–1247.

[4] A. Behera and G. K. Panda, *On the square roots of triangular numbers*, Fibonacci Quart. 37 (1999) 98–105.

[5] R. Finkelstein, *The house problem*, Amer. Math. Monthly **72** (1965), 1082–1088.

[6] T. Komatsu, *Higher-order convolution identities for Cauchy numbers of the second kind*, Proc. Jangjeon Math. Soc. **18** (2015), 369–383.

[7] T. Komatsu, *Higher-order convolution identities for Cauchy numbers*, Tokyo J. Math. **39** (2016). 15 pages.

[8] T. Komatsu, Z. Masakova and E. Pelantova, *Higher-order identities for Fibonacci numbers*, Fibonacci Quart. **52**, no.5 (2014), 150–163.

[9] T. Komatsu and L. Szalay, *Balancing with binomial coefficients*, Intern. J. Number Theory **10** (2014), 1729–1742.

[10] T. Komatsu and Y. Simsek, *Third and higher order convolution identities for Cauchy numbers*, Filomat **30** (2016), 1053–1060.

[11] K. Liptai, *Fibonacci Balancing numbers*, Fibonacci Quart. **42** (2004), 330–340.

[12] K. Liptai, F. Luca, Á. Pintér and L. Szalay, *Generalized balancing numbers*, Indag. Math. (N.S.) **20** (2009), 87–100.

[13] G. K. Panda, *Some fascinating properties of balancing numbers*, In Proc. of Eleventh Internat. Conference on Fibonacci Numbers and Their Applications, Cong. Numerantium **194** (2009), 185–189.

[14] B. K. Patel and P. K. Ray, *The period, rank and order of the sequence of balancing numbers modulo m*, Math. Rep. (Bucur.) **18**, No.3 (2016), Article No.9.

[15] P. K. Ray, *Some congruences for balancing and Lucas-balancing numbers and their applications*, Integers, **14** (2014), #A8.